

Understanding Limits

In algebra, we are often required to calculate an exact value of a function $y=f(x)$. When determining a specific y -value of a function, we simply need the corresponding x -value.

However, what if the function isn't just a simple curve? For example, we will be examining functions with holes, functions that shoot up toward infinity, functions that oscillate, etc. To describe the behaviour of the y -values of such functions, we can use a *limit*.

Intuitive Definition of a Limit

The limit of a function describes how a function behaves *near* a specific point, but not *at* that point. If the values of $y=f(x)$ get closer and closer to one number, L , as we take values of x very close to (but not equal to) a number, c , then we say "**The limit of $f(x)$, as x approaches c , is L** " and we *write*:

$$\lim_{x \rightarrow c} f(x) = L$$

It is important to note that the value of the *limit* at $x = c$ does *not* depend on the value of the *function* at $x = c$.

- $f(c)$ is a single number that describes the value of $f(x)$ *at* point $x = c$.
- $\lim_{x \rightarrow c} f(x)$ is a single number that describes the behavior of $f(x)$ *near, but not at*, the point $x = c$.

Using Tables to Find Limits

One way to approximate a limit is to use a table of values. Choose x -values close to c from both the left and the right. If the y -values approach a distinct value, L , as x approaches c , then the limit is L .

Example 1:

Determine the value of $\lim_{x \rightarrow 5} \frac{x^2 - 3x - 10}{x - 5}$

Solution:

1. Create a table of values for the function $f(x) = \frac{x^2 - 3x - 10}{x - 5}$ and fill in values of x that get close to 5 from both sides. Then, calculate the corresponding $f(x)$ values.

x	4	4.5	4.9	4.95	5	5.05	5.1	5.5	6
$f(x)$									

2. Examine the table to determine if the $f(x)$ values approach the same number from the left and the right.

$$\lim_{x \rightarrow 5} \frac{x^2 - 3x - 10}{x - 5} = \underline{\hspace{2cm}}$$

* Note that we cannot find the limit by substituting $x = 5$ into the function since this results in division by zero.

Using Graphs to Find Limits

If we have the graph of a function $f(x)$ near $x = c$, then it is usually easy to determine $\lim_{x \rightarrow c} f(x)$.

Example 2:

Use the graph of the function to the right to determine the following limits.

- a. $\lim_{x \rightarrow 4} f(x)$ b. $\lim_{x \rightarrow 2} f(x)$

Solution:

- a. As the x -values get closer to 4 from the left and from the right (We will learn more about Left- & Right-Hand Limits later), the y -values get closer to ____.


Note that $f(4) = \underline{\hspace{2cm}}$, but the limit of $f(x)$ as x approaches 4 is ____.

$$\lim_{x \rightarrow 4} f(x) = \underline{\hspace{2cm}}$$

- b. As the x -values get closer to 2 from the left and from the right, the y -values get closer to ____.

Note that $f(2) = \underline{\hspace{2cm}}$. In this case, the function value and the limit value are the same.

$$\lim_{x \rightarrow 2} f(x) = \underline{\hspace{2cm}}$$



Example 3:

Use the graph of the function to the right to determine the following limits.

- a. $\lim_{x \rightarrow a} f(x)$ b. $\lim_{x \rightarrow b} f(x)$ c. $\lim_{x \rightarrow c} f(x)$

Solution:

- a. As the x -values get closer to a from either side, the y -values get closer to ____.

Note that the function doesn't exist at a , but the limit does.

$$\lim_{x \rightarrow a} f(x) = \underline{\hspace{2cm}}$$

- b. As the x -values get closer to b from the left, the y -values get closer to ____.
As the x -values get closer to b from the right, the y -values get closer to ____.


Since the function approaches different values from each side, the limit *does not exist*.

$$\lim_{x \rightarrow b} f(x) \underline{\hspace{2cm}}$$

- c. As the x -values get closer to c from either side, the y -values get closer to ____.

Since the function approaches infinity from each side, the limit *does not exist*.

$$\lim_{x \rightarrow c} f(x) \underline{\hspace{2cm}}$$



Calculating Limits Algebraically

Limit Rules

If n is a positive integer, k is a constant, and $f(x)$ and $g(x)$ are functions that have limits at $x = c$, then the following rules hold:

1. Constant Rule

$$\lim_{x \rightarrow c} k = k$$

2. Identity Rule

$$\lim_{x \rightarrow c} x = c$$

3. Constant Multiple Rule

$$\lim_{x \rightarrow c} kf(x) = k \lim_{x \rightarrow c} f(x)$$

4. Sum Rule

$$\lim_{x \rightarrow c} [f(x) + g(x)] = \lim_{x \rightarrow c} f(x) + \lim_{x \rightarrow c} g(x)$$

5. Difference Rule

$$\lim_{x \rightarrow c} [f(x) - g(x)] = \lim_{x \rightarrow c} f(x) - \lim_{x \rightarrow c} g(x)$$

6. Product Rule

$$\lim_{x \rightarrow c} [f(x) \cdot g(x)] = \lim_{x \rightarrow c} f(x) \cdot \lim_{x \rightarrow c} g(x)$$

7. Quotient Rule

$$\lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow c} f(x)}{\lim_{x \rightarrow c} g(x)}, \text{ if } \lim_{x \rightarrow c} g(x) \neq 0$$

8. Power Rule

$$\lim_{x \rightarrow c} [f(x)]^n = \left[\lim_{x \rightarrow c} f(x) \right]^n$$

9. n^{th} Root Rule

$$\lim_{x \rightarrow c} \sqrt[n]{f(x)} = \sqrt[n]{\lim_{x \rightarrow c} f(x)}, \text{ if } \lim_{x \rightarrow c} f(x) > 0$$

Example 4: Finding a Limit Using Limit Rules


Find $\lim_{x \rightarrow -2} 3$

Solution:

Use the Constant Rule for limits.

$$\lim_{x \rightarrow -2} 3 = \underline{\hspace{2cm}}$$

Look at the graph of the function $y = 3$ to see why the limit of the constant function as x approaches 3 is the same as the constant value. Notice that for this function, the limit as x approaches *any* value is 3.



Example 5: Finding a Limit Using Limit RulesFind $\lim_{x \rightarrow 3} (2x^2 - 4x - 1)$ **Solution:**

Use the Difference Rule.

$$\lim_{x \rightarrow 3} (2x^2 - 4x - 1) = \lim_{x \rightarrow 3} 2x^2 - \lim_{x \rightarrow 3} 4x - \lim_{x \rightarrow 3} 1$$

Use the Constant Multiple Rule.

$$= 2 \lim_{x \rightarrow 3} x^2 - 4 \lim_{x \rightarrow 3} x - \lim_{x \rightarrow 3} 1$$

Use the Power Rule.

$$= 2 \left(\lim_{x \rightarrow 3} x \right)^2 - 4 \lim_{x \rightarrow 3} x - \lim_{x \rightarrow 3} 1$$

Use the Constant Rule.

$$= 2 \left(\lim_{x \rightarrow 3} x \right)^2 - 4 \lim_{x \rightarrow 3} x - 1$$

Substitute.


$$= 2(3)^2 - 4(3) - 1$$

Simplify.

$$= 5$$

Note that the limit as x approaches 3 is the same as the function value at $x = 3$.

This is always the case for limits of polynomial functions.

We can use *direct substitution* to evaluate all polynomial limits.**Example 6: Finding a Limit by Direct Substitution**

a. Find $\lim_{x \rightarrow -1} (-x^4 + 5x^3 + 3x^2 - 1)$ b. Find $\lim_{x \rightarrow 8} \frac{x^2 - 2x - 3}{x + 2}$

Solution:

a. Use direct substitution.

$$\lim_{x \rightarrow -1} (-x^4 + 5x^3 + 3x^2 - 1) = -(-1)^4 + 5(-1)^3 + 3(-1)^2 - 1$$

$$= \underline{\hspace{2cm}}$$

$$= \underline{\hspace{1cm}}$$

b. Use direct substitution.

$$\lim_{x \rightarrow 8} \frac{x^2 - 2x - 3}{x + 2} = \frac{8^2 - 2(8) - 3}{8 + 2}$$

$$=$$

$$=$$

Example 7: Finding a Limit by Factoring & Simplifying First

a. Find $\lim_{x \rightarrow 1} \frac{2x^2 - x - 1}{x - 1}$

b. Find $\lim_{x \rightarrow 2} \frac{x^3 - 8}{x - 2}$

Solution:

a. If we try to evaluate $\lim_{x \rightarrow 1} \frac{2x^2 - x - 1}{x - 1}$ by using direct substitution, we get $\frac{2(1)^2 - 1 - 1}{1 - 1} = \frac{0}{0}$.

So, factor and simplify first, *then* use substitution.

$$\lim_{x \rightarrow 1} \frac{2x^2 - x - 1}{x - 1} =$$

b. If we try to evaluate $\lim_{x \rightarrow 2} \frac{x^3 - 8}{x - 2}$ by using direct substitution, we get $\frac{(2)^3 - 8}{2 - 2} = \frac{0}{0}$.

So, factor and simplify first, *then* use substitution.

Example 8: Finding a Limit by Using the Conjugate Technique

a. Find $\lim_{x \rightarrow 4} \frac{x-4}{\sqrt{x}-2}$ b. Find $\lim_{x \rightarrow -3} \frac{2-\sqrt{x^2-5}}{x+3}$

Solution:

a. Direct substitution yields $\frac{0}{0}$. So, multiply by the conjugate form of 1.


b. Direct substitution yields $\frac{0}{0}$. So, multiply by the conjugate form of 1.

$$\lim_{x \rightarrow -3} \frac{2-\sqrt{x^2-5}}{x+3}$$


Practice:

1. Determine the following limits graphically.


a. $\lim_{x \rightarrow 1} f(x)$




b. $\lim_{x \rightarrow 1} f(x)$




c. $\lim_{x \rightarrow -1} f(x)$




d. $\lim_{x \rightarrow 3} f(x)$



e. $\lim_{x \rightarrow \frac{\pi}{6}} f(x)$



f. $\lim_{x \rightarrow 4} f(x)$



2. Use the *Limit Rules* to calculate $\lim_{x \rightarrow 3} \frac{(4x+2)^2}{\sqrt[3]{1-x^2}}$.

3. Evaluate the following limits.

a) $\lim_{x \rightarrow 4} \frac{x-4}{x^3-64}$

j) $\lim_{x \rightarrow -8} \frac{x+8}{\sqrt[3]{x}+2}$

s) $\lim_{x \rightarrow 5} \frac{2x}{x-5}$

b) $\lim_{x \rightarrow 7} \frac{\sqrt[5]{3-5x}}{(x-5)^3}$

k) $\lim_{x \rightarrow -1} \frac{8}{(3+x)\sqrt{3-x}}$

t) $\lim_{x \rightarrow 0} \frac{x^3}{2x^3+3x^4}$

c) $\lim_{x \rightarrow -3} \frac{x+3}{3-\sqrt{x+12}}$

l) $\lim_{x \rightarrow 9} \frac{x-9}{3-\sqrt{x}}$

u) $\lim_{x \rightarrow 2} \frac{x^2-2x}{2x^2-7x+6}$

d) $\lim_{x \rightarrow -3} \sqrt[3]{\frac{x-4}{6x^2+2}}$

m) $\lim_{x \rightarrow 0} \frac{(3+x)^3-3^3}{(3+x)^2-3^2}$

v) $\lim_{x \rightarrow \pi} \frac{1+\sin x}{\cos x}$

e) $\lim_{x \rightarrow 3} \frac{\frac{1}{x^2}-\frac{1}{9}}{x-3}$

n) $\lim_{x \rightarrow 9} \frac{x^2-81}{3-\sqrt{x}}$

w) $\lim_{x \rightarrow \frac{\pi}{4}} ((\cos x) + (\sin x))$

f) $\lim_{h \rightarrow 0} \frac{(-3+h)^2-9}{h}$

o) $\lim_{x \rightarrow 16} \frac{2\sqrt{x}+x^{3/2}}{\sqrt[4]{x}+5}$

x) $\lim_{x \rightarrow \frac{3\pi}{2}} \frac{\sin x}{\cos x}$

g) $\lim_{h \rightarrow 0} \frac{(2+h)^3-8}{h}$

p) $\lim_{x \rightarrow 2} \frac{x^2-4}{3x^2-7x+2}$

y) $\lim_{x \rightarrow 4} \frac{2x^3-128}{\sqrt{x}-2}$

h) $\lim_{x \rightarrow 0} \frac{\frac{1}{1+x}-1}{x}$

q) $\lim_{x \rightarrow 7} \frac{\sqrt{x+2}-3}{x-7}$

i) $\lim_{x \rightarrow -1} \frac{2x+3}{3x+2}$

r) $\lim_{x \rightarrow 4} \left(\frac{8}{x^2-16} - \frac{1}{x-4} \right)$

Answers:

1. a. 2 b. 0 c. 3 d. DNE e. 2 f. 6

2.

$$\begin{aligned} \lim_{x \rightarrow 3} \frac{(4x+2)^2}{\sqrt[3]{1-x^2}} &= \frac{\lim_{x \rightarrow 3} (4x+2)^2}{\lim_{x \rightarrow 3} \sqrt[3]{1-x^2}} = \frac{\left(\lim_{x \rightarrow 3} (4x+2) \right)^2}{\sqrt[3]{\lim_{x \rightarrow 3} (1-x^2)}} = \frac{\left(\lim_{x \rightarrow 3} (4x) + \lim_{x \rightarrow 3} 2 \right)^2}{\sqrt[3]{\lim_{x \rightarrow 3} 1 - \lim_{x \rightarrow 3} (x^2)}} = \frac{\left(4 \lim_{x \rightarrow 3} x + \lim_{x \rightarrow 3} 2 \right)^2}{\sqrt[3]{\lim_{x \rightarrow 3} 1 - \left(\lim_{x \rightarrow 3} x \right)^2}} \\ &= \frac{(4(3)+2)^2}{\sqrt[3]{1-(3)^2}} \\ &= \frac{196}{\sqrt[3]{-8}} = \frac{196}{-2} = -98 \end{aligned}$$

3. a) $\frac{1}{48}$ b) $-\frac{1}{4}$ c) -6 d) $-\frac{1}{2}$ e) $-\frac{2}{27}$ f) -6 g) 12 h) -1

i) -1 j) 12 k) 2 l) -6 m) $\frac{9}{2}$ n) -108 o) $\frac{72}{7}$ p) $\frac{4}{5}$ q) $\frac{1}{6}$

r) $-\frac{1}{8}$ s) DNE t) $\frac{1}{2}$ u) 2 v) -1 w) $\sqrt{2}$ x) DNE y) 384