

# Infinite Limits

There are many examples of functions that increase and/or decrease without bound. These functions have limit behaviour that does not exist. If the y-values increase or decrease without bound as  $x$  approaches  $c$ , we can write  $\lim_{x \rightarrow c} f(x) = +\infty$  or  $\lim_{x \rightarrow c} f(x) = -\infty$ , respectively. It is important to note that infinity is a behaviour, not a value. Stating that the limit of a given function is  $+\infty$  or  $-\infty$  is simply a *specific* way of saying that the limit does not exist.


## Example 1:

Find  $\lim_{x \rightarrow a} f(x)$

### Solution:

By looking at the graph of  $y=f(x)$ , we can see that the y-values *increase* without bound as  $x$  approaches  $a$  from the left and from the right.

$$\lim_{x \rightarrow a} f(x) = \underline{\hspace{2cm}}$$




## Example 2:

Find  $\lim_{x \rightarrow a} f(x)$

### Solution:

By looking at the graph of  $y=f(x)$ , we can see that the y-values *decrease* without bound as  $x$  approaches  $a$  from the left and from the right.

$$\lim_{x \rightarrow a} f(x) = \underline{\hspace{2cm}}$$



## Example 3:

- a. Find  $\lim_{x \rightarrow a^-} f(x)$       b. Find  $\lim_{x \rightarrow a^+} f(x)$


### Solution:

- a. By looking at the graph of  $y=f(x)$ , we can see that the y-values *decrease* without bound as  $x$  approaches  $a$  from the *left*.

$$\lim_{x \rightarrow a^-} f(x) = \underline{\hspace{2cm}}$$

- b. By looking at the graph of  $y=f(x)$ , we can see that the y-values *increase* without bound as  $x$  approaches  $a$  from the *right*.

$$\lim_{x \rightarrow a^+} f(x) = \underline{\hspace{2cm}}$$



# Vertical Asymptotes and Points of Discontinuity

To find these features of a given function, first factor the numerator and denominator. There will be either a vertical asymptote or a point of discontinuity at any x-value that would cause the denominator to equal zero.

Next, cancel like factors. If this eliminates the division by zero, then the function has a hole in the graph at the corresponding x-value. If canceling like factors does not eliminate the division by zero, then the function has a vertical asymptote at the corresponding x-value.

Once we have found the vertical asymptotes and points of discontinuity of a function, we can use limits to determine the behaviour of the function on each side of an asymptote and the y-coordinate of a point of discontinuity.

Functions that have *vertical asymptotes* approach positive or negative *infinity* as the x-values approach an asymptote from the left- and/or right-hand sides.

Functions that have *points of discontinuity* (holes) approach a *finite* value as the x-values approach a point of discontinuity from both sides.

## Example 4:

Use factoring and simplifying to determine if there are any vertical asymptotes or points of discontinuity for the

$$\text{function } f(x) = \frac{x^2 - 3x + 2}{x^2 - 4x + 3}.$$

### Solution:

$$f(x) = \frac{x^2 - 3x + 2}{x^2 - 4x + 3} =$$

Since the factor  $(x-1)$  cancels out of the denominator, we know there is a \_\_\_\_\_ at  $x = \underline{\hspace{1cm}}$ .

Since the factor  $(x-3)$  doesn't cancel out of the denominator, we know there is a \_\_\_\_\_ at  $x = \underline{\hspace{1cm}}$ .

## Example 5:

As discovered in the previous example, the function  $f(x) = \frac{x^2 - 3x + 2}{x^2 - 4x + 3}$  has a point of discontinuity at  $x = 1$ .

Find the limit of  $f(x)$  as  $x$  approaches 1 in order to determine the y-coordinate of this point.

### Solution:

$$\lim_{x \rightarrow 1} \frac{x^2 - 3x + 2}{x^2 - 4x + 3} = \lim_{x \rightarrow 1} \frac{(x-2)(x-1)}{(x-3)(x-1)} = \lim_{x \rightarrow 1} \frac{(x-2)}{(x-3)} =$$

Therefore, the coordinates of the P.O.D. are \_\_\_\_\_.

**Example 6:**

As discovered in example 4, the function  $f(x) = \frac{x^2 - 3x + 2}{x^2 - 4x + 3}$  has a vertical asymptote at  $x = 3$ . Find the limit of  $f(x)$  as  $x$  approaches 3 from the left and from the right in order to determine the behaviour of the function on each side of the asymptote.

**Solution:**

Substitute an  $x$ -value very close to 3, approaching from the left ( $x < 3$ ), to calculate the following limit:

$$\lim_{x \rightarrow 3^-} \frac{x^2 - 3x + 2}{x^2 - 4x + 3} = \lim_{x \rightarrow 3^-} \frac{(x-2)(x-1)}{(x-3)(x-1)} = \lim_{x \rightarrow 3^-} \frac{(x-2)}{(x-3)} =$$


Therefore, the function \_\_\_\_\_ without bound as  $x$  approaches 3 from the left.

Substitute an  $x$ -value very close to 3, approaching from the right ( $x > 3$ ), to calculate the following limit:

$$\lim_{x \rightarrow 3^+} \frac{x^2 - 3x + 2}{x^2 - 4x + 3} = \lim_{x \rightarrow 3^+} \frac{(x-2)(x-1)}{(x-3)(x-1)} = \lim_{x \rightarrow 3^+} \frac{(x-2)}{(x-3)} =$$

Therefore, the function \_\_\_\_\_ without bound as  $x$  approaches 3 from the right.

Note that the results of the previous examples are consistent with the features on the graph of  $f(x)$ :



# Limits and End Behaviour: Horizontal Asymptotes

In order to determine the end behaviour of a function, we can evaluate limits at infinity. That is, we calculate a limit (algebraically, graphically, or numerically) as  $x$  approaches positive or negative infinity.


Using the result of the limit calculation, we can determine whether or not the graph of the function has a horizontal asymptote.

When direct substitution into the function produces the indeterminate form  $\frac{\pm\infty}{\pm\infty}$  or  $\infty - \infty$ , we must be careful.

These are cases which often have *finite* limits that can be determined through some algebraic manipulation.


For any function, there are three possibilities for the *limit at infinity*:

## Limit Does Not Exist




No horizontal asymptote

## Limit is Zero



Horizontal asymptote at  $y = 0$

## Limit is a Non-Zero Number



Horizontal asymptote at  $y = \text{limit}$   
(In this case,  $y = 2$ )

## Example 7: Determining a Limit at Infinity

Determine the end-behaviour of the function  $f(x) = \frac{10x^2 + 7x}{3x^2 - 5x + 1}$  by calculating  $\lim_{x \rightarrow \infty} \frac{10x^2 + 7x}{3x^2 - 5x + 1}$ .

**Solution:**

Direct substitution gives  $\frac{\infty}{\infty}$ .

Divide the numerator and denominator by the highest power of  $x$ , simplify, and then substitute.

$$\lim_{x \rightarrow \infty} \frac{10x^2 + 7x}{3x^2 - 5x + 1} =$$

Therefore, there is a horizontal asymptote for the function  $f(x) = \frac{10x^2 + 7x}{3x^2 - 5x + 1}$  at \_\_\_\_\_.

**Example 8: Determining a Limit at Infinity**

Determine the end-behaviour of the function  $f(x) = \frac{4x^3 - 6x^2 + 8}{7x^5 + 6x^2 - 9x + 2}$  by calculating  $\lim_{x \rightarrow -\infty} \frac{4x^3 - 6x^2 + 8}{7x^5 + 6x^2 - 9x + 2}$ .

**Solution:**

Direct substitution gives  $\frac{\infty}{\infty}$ .

Divide the numerator and denominator by the highest power of  $x$ , simplify, and then substitute.

$$\lim_{x \rightarrow -\infty} \frac{4x^3 - 6x^2 + 8}{7x^5 + 6x^2 - 9x + 2} =$$

Therefore, there is a horizontal asymptote for the function  $f(x) = \frac{4x^3 - 6x^2 + 8}{7x^5 + 6x^2 - 9x + 2}$  at \_\_\_\_\_.

**Example 9: Determining a Limit at Infinity**

Determine the end-behaviour of the function  $f(x) = \frac{3 - 4x^2}{5x - 1}$  by calculating  $\lim_{x \rightarrow \pm\infty} \frac{3 - 4x^2}{5x - 1}$ .

**Solution:**

Direct substitution gives  $\frac{\infty}{\infty}$  or  $\frac{-\infty}{\infty}$ .

Divide the numerator and denominator by the highest power of  $x$ , simplify, and then substitute.

$$\lim_{x \rightarrow \pm\infty} \frac{3 - 4x^2}{5x - 1} =$$

Therefore, there is *no* horizontal asymptote for the function  $f(x) = \frac{3 - 4x^2}{5x - 1}$ .


To determine whether the function *increases* or *decreases* without bound, we can determine the limits at infinity separately (That is, the limit as  $x$  approaches *negative* infinity *and* the limit as  $x$  approaches *positive* infinity).

Substitute any representative  $x$ -value (very small, tending toward negative infinity, and then very large, tending toward positive infinity), into the *leading terms* of the numerator and denominator to determine if the overall limit will be *positive* or *negative* infinity.

$$\lim_{x \rightarrow -\infty} \frac{3 - 4x^2}{5x - 1} = \frac{-4(\underline{\hspace{2cm}})^2}{5(\underline{\hspace{2cm}})} =$$

$$\lim_{x \rightarrow +\infty} \frac{3 - 4x^2}{5x - 1} = \frac{-4(\underline{\hspace{2cm}})^2}{5(\underline{\hspace{2cm}})} =$$

Note that these limits reflect the end behaviour of the graph of  $f(x) = \frac{3 - 4x^2}{5x - 1}$ :



**Example 10: Limits at Infinity Involving Radicals**

Determine  $\lim_{x \rightarrow +\infty} \frac{\sqrt{2x^3 - x}}{3x^2 - 6}$

**Solution:**

Direct substitution gives  $\frac{\infty}{\infty}$ .

Divide the numerator and denominator by the highest power of  $x$ . Recall that, for example,  $x^2 = \sqrt{x^4}$ .

$$\lim_{x \rightarrow +\infty} \frac{\sqrt{2x^3 - x}}{3x^2 - 6} =$$

**Example 11: Limits at Infinity Involving Radicals**

Determine  $\lim_{x \rightarrow +\infty} (\sqrt{x^2 - 4x} - x)$

**Solution:**

Direct substitution gives  $\infty - \infty$ .

Multiply by the conjugate form of 1.

$$\lim_{x \rightarrow +\infty} (\sqrt{x^2 - 4x} - x) =$$

## Practice:

1. Explain in your own words the meaning of each of the following:

a)  $\lim_{x \rightarrow -\infty} f(x) = 6$

b)  $\lim_{x \rightarrow \infty} f(x) = -9$

c)  $\lim_{x \rightarrow 4^+} f(x) = \infty$

d)  $\lim_{x \rightarrow 6^-} f(x) = -\infty$

2. Can the graph of  $y = f(x)$  intersect the following? Explain.

a) vertical asymptote

b) horizontal asymptote

3. For the function whose graph is shown below, determine the following:


a)  $\lim_{x \rightarrow \infty} f(x)$

b)  $\lim_{x \rightarrow -\infty} f(x)$

c)  $\lim_{x \rightarrow -3^-} f(x)$

d)  $\lim_{x \rightarrow -3^+} f(x)$

e) State the equations of the vertical and horizontal asymptotes.



4. Evaluate  $\lim_{x \rightarrow \infty} \frac{x^2}{2^x}$  by using a table of values.

5. Evaluate the following limits.

a)  $\lim_{x \rightarrow \infty} \frac{5x^4 - 7x^3 + 7x^2 - 1}{3x^4 + 2x^3}$

b)  $\lim_{x \rightarrow -\infty} \frac{x^5 - x^2}{x^3 - 2x}$

c)  $\lim_{x \rightarrow \infty} \frac{9x^2 - x + 8}{2x^4 + x^3 - 7}$

d)  $\lim_{x \rightarrow -\infty} \frac{12x^2 - 6x^3 + 5x^4 + 9x^5}{3x^5 + 2x^4 - 4x^3 + 2x}$

e)  $\lim_{x \rightarrow \infty} \frac{5x^3 - 4x^2 - 5x}{4x^3 + 3x}$

f)  $\lim_{x \rightarrow \infty} \left[ \left(\frac{1}{8}\right)^x + \frac{x^3 - 4x^2 - 5x}{4x^3 + 3x} - 7 \right]$

g)  $\lim_{x \rightarrow \infty} \left[ \left(\frac{7}{3}\right)^{-x} + \frac{4x^2 - 5x}{2x^2 + 1} - 9 \right]$

h)  $\lim_{x \rightarrow -\infty} \left(\frac{1}{5}\right)^{2x}$

i)  $\lim_{x \rightarrow \infty} \frac{(-2x^2 - 3)(x + 1)}{2 - 5x^3}$

j)  $\lim_{x \rightarrow -\infty} \left[ 12 - \left(\frac{6}{5}\right)^x \right]$

k)  $\lim_{x \rightarrow \infty} \frac{10x^2}{\sqrt{4x^4 + 1}}$

l)  $\lim_{x \rightarrow \infty} \frac{2x}{\sqrt{x^2 - 2} - 3x}$

m)  $\lim_{x \rightarrow \infty} \frac{4 - \frac{3}{x}}{5x^2 + 1}$

n)  $\lim_{x \rightarrow \infty} \left[ \frac{6x}{2x - 1} - \frac{x + 5}{3x - 4} \right]$

o)  $\lim_{x \rightarrow \infty} [x - \sqrt{x^2 + 1}]$

p)  $\lim_{x \rightarrow -\infty} \frac{|x - 8|}{x - 8}$

q)  $\lim_{x \rightarrow \infty} [2x - \sqrt{4x^2 + 6x}]$

r)  $\lim_{x \rightarrow -\infty} (x - 3)^2(x + 1)^5$

s)  $\lim_{x \rightarrow -\infty} (2x - 3)^3(x + 1)^3$

6. Consider each of the following functions:

$$\text{i. } f(x) = \frac{x^2 + 8x - 20}{2x^2 + x - 6}$$

$$\text{ii. } f(x) = \frac{-2x + 4}{x^3 + 4x^2 - 3x - 18}$$

$$\text{iii. } f(x) = \frac{x^4 - 2x^3 - 63x^2}{x^2 - 10x + 16}$$

$$\text{iv. } f(x) = \frac{10x^3 - 18x}{x^3 - x^2 - 2x}$$

$$\text{v. } f(x) = \frac{9 - x^2}{16 - 2x^3}$$

$$\text{vi. } f(x) = \frac{x^3 + 4x^2 + 3x}{x^2 + 8x + 15}$$


- determine if there are any vertical asymptotes and/or points of discontinuity, and for which values of  $x$  these occur.
- use limits** to determine the behavior of  $f(x)$  on each side of any vertical asymptote that exists.
- use limits** to determine the coordinates of any point of discontinuity that exists.
- use limits** to determine whether or not a horizontal asymptote exists and, if so, what its equation is.


## Answers:


- As  $x$  approaches negative infinity, the value of the function approaches **6**.
  - As  $x$  approaches positive infinity, the value of the function approaches **-9**.
  - As  $x$  approaches **4** from the right hand side, the value of the function gets increasingly large, approaching positive infinity.
  - As  $x$  approaches **6** from the left hand side, the value of the function becomes more negative, approaching negative infinity.
- No, the function is undefined at a vertical asymptote.
  - Yes, the graph of a function can cross a horizontal asymptote. A horizontal asymptote describes the *end-behaviour* of a function - its tendency to approach a particular  $y$ -value as  $x$  approaches positive or negative infinity.
- 4**
  - 1**
  - $\infty$
  - $-\infty$
  - HA :  $y = 1$  and  $y = 4$**   
**VA :  $x = -3$**
- $$\lim_{x \rightarrow \infty} \frac{x^2}{2^x} = 0$$
- $\frac{5}{3}$
  - $\infty$
  - 0**
  - 3**
  - $\frac{5}{4}$
  - $-\frac{27}{4}$
  - 7**
  - $\infty$
  - $\frac{2}{5}$
  - 12**
  - 5**
  - 1**
  - 0**
  - $\frac{8}{3}$
  - 0**
  - 1**
  - $-\frac{3}{2}$
  - $-\infty$
  - $\infty$





6.


<p>i)</p> $f(x) = \frac{x^2 + 8x - 20}{2x^2 + x - 6}$	<p>VA  <math>x = 3/2</math>  <math>x = -2</math></p>	$\lim_{x \rightarrow -2^-} f(x) = -\infty$ $\lim_{x \rightarrow -2^+} f(x) = \infty$ $\lim_{x \rightarrow \frac{3}{2}^-} f(x) = \infty$ $\lim_{x \rightarrow \frac{3}{2}^+} f(x) = -\infty$	<p>No POD</p>	$\lim_{x \rightarrow \infty} f(x) = \frac{1}{2}$ <p>HA  <math>y = \frac{1}{2}</math></p>	
---	--	---	---------------	--	---

<p>ii)</p> $f(x) = \frac{-2x + 4}{x^3 + 4x^2 - 3x - 18}$	<p>VA  <math>x = -3</math>                  POD                  at  <math>x = 2</math></p>	$\lim_{x \rightarrow -3^-} f(x) = -\infty$ $\lim_{x \rightarrow -3^+} f(x) = -\infty$	<p>POD                  (2, -0.08)</p>	$\lim_{x \rightarrow \infty} f(x) = 0$ <p>HA  <math>y = 0</math></p>	
--	---	---	--	--	--

<p>iii)</p> $f(x) = \frac{x^4 - 2x^3 - 63x^2}{x^2 - 10x + 16}$	<p>VA  <math>x = 2</math>  <math>x = 8</math></p>	$\lim_{x \rightarrow 2^-} f(x) = -\infty$ $\lim_{x \rightarrow 2^+} f(x) = \infty$ $\lim_{x \rightarrow 8^-} f(x) = \infty$ $\lim_{x \rightarrow 8^+} f(x) = -\infty$	<p>No POD</p>	$\lim_{x \rightarrow \infty} f(x) = DNE$ <p>No HA</p>	
--	---	---	---------------	---	---

<p>iv)</p> $f(x) = \frac{10x^3 - 18x}{x^3 - x^2 - 2x}$	<p>VA  <math>x = -1</math>  <math>x = 2</math></p>	<p><math>\lim_{x \rightarrow -1^-} f(x) = -\infty</math>  <math>\lim_{x \rightarrow -1^+} f(x) = \infty</math>  <math>\lim_{x \rightarrow 2^-} f(x) = -\infty</math>  <math>\lim_{x \rightarrow 2^+} f(x) = \infty</math></p>	<p>POD  <math>(0, 9)</math></p>	<p><math>\lim_{x \rightarrow \infty} f(x) = 10</math>                      HA  <math>y = 10</math></p>	
--	--	---	-------------------------------------	--	---

<p>v)</p> $f(x) = \frac{9 - x^2}{16 - 2x^3}$	<p>VA  <math>x = 2</math></p>	<p><math>\lim_{x \rightarrow 2^-} f(x) = \infty</math>  <math>\lim_{x \rightarrow 2^+} f(x) = -\infty</math></p>	<p>No POD</p>	<p><math>\lim_{x \rightarrow \infty} f(x) = 0</math>                      HA  <math>y = 0</math></p>	
--	-----------------------------------	--	---------------	--	--

<p>vi)</p> $f(x) = \frac{x^3 + 4x^2 + 3x}{x^2 + 8x + 15}$	<p>VA  <math>x = -5</math>                      POD at  <math>x = -3</math></p>	<p><math>\lim_{x \rightarrow -5^-} f(x) = -\infty</math>  <math>\lim_{x \rightarrow -5^+} f(x) = \infty</math></p>	<p>POD  <math>(-3, 3)</math></p>	<p><math>\lim_{x \rightarrow \infty} f(x) = DNE</math>                      No HA</p>	
---	---	--	--------------------------------------	---	---